

- Facilities:
- Ascension St. Vincent's Riverside
 - Ascension St. Vincent's Southside
 - Ascension St. Vincent's Clay County
 - Ascension St. Johns County

AFFIX PATIENT LABEL

RAD Gastrostomy/Gastrojejunostomy Tube Post-Placement Non Categorized

Last Revision Date

- 2023/11

Vital Signs

Vital Signs

- T;N q15min x 4, q30min x 2, then per protocol.

Procedural Site Check

- once following procedure.

Call Provider for the following:

- Contact radiologist for any unstable vital signs, changes in patient's symptoms, or procedural site complications.

Activity

Activity

- Bedrest for 2 hrs, then resume pre procedure activity.

Diet

Diets

- Resume diet after procedure.

Patient Care

Call Provider for the following:

- T;N Contact Interventional Radiology if catheter is malfunctioning, poorly functioning, leaking or broken.

Gastric Tube

- Feeding tube may be used starting 2 hours after placement. Flush tube with 50 mL sterile water after each use.

Gastrojejunostomy (2 lumen) Tube

- Feeding tube may be used starting 2 hours after placement. Flush tube with 50 mL sterile water after each use.

Medications

Analgesics

LINKED CONTENT: Adult Multimodal Acute Pain with TAG NATL

Antiemetics

ondansetron

- 4 mg inj IV slow push once PRN nausea

promethazine

- 12.5 mg inj IV slow push once PRN nausea If zofran also ordered, use zofran first.

- 6.25 mg inj IV slow push once PRN nausea For age > or = 65 years For age < 65 years order 12.5mg order. Decrease to 6.25mg for patients age >/= 65 years. If Zofran ordered, use Zofran first.

Nursing Signature _____ Date/Time _____

Physician Signature _____ Date/Time _____

