

CT Abdomen/Pelvis Routine

Updated
5/2/2024

Indications - abdominal pain, nausea, vomiting, constipation, diarrhea, weight loss, fever, metastatic workup, etc.

GENERAL SCAN NOTES

Move the patient's arms over his/her head if possible. Remove any metal from the imaging field of view.

Topogram - lung bases through pubic symphysis (obtained during end inspiration).

Craniocaudal scan coverage - lung bases through pubic symphysis (obtained during end inspiration).

Adjust FOV (field of view) on topogram to smallest without cropping anatomy.

IV Contrast: 100 mL Omnipaque-300, inject at 2.5 mL/sec.

70 secs scan delay for most indications.

90 secs scan delay for renal indications like pyelonephritis, renal abscess, renal infarct or renal vein thrombosis.

Do not do delayed/excretory phase unless asked to do so.

Oral Contrast: per protocol.

For **GE scanners**, it is essential for the 1st recon thickness on the scanner to match the 1st recon thickness in this protocol book for the prescribed Noise Index to be valid. The 1st recon should generally be the thickest recon in the protocol.

SIEMENS PARAMETERS & RECONS

	Scan Mode	kV	mAs	Care Dose	Care kV & Lvl	Pitch	Acq	Coll	Rot Time	Scan Time
Sensation 16	spiral	120	200	on	NA	0.80	16	1.5	0.5	13.0
Go Up 32	spiral	130	92	on	on 145	0.80	32	0.7	0.8	22.3
Sensation 64	spiral	120	200	on	NA	1.20	24	1.2	0.5	7.2
Definition 64	spiral	120	180	on	on	0.60	64	0.6	0.5	21.7
Go Top 64	spiral	120	112	on	on 145	0.80	64	0.6	0.5	8.1
Drive 128	spiral	120	147	on	on	0.60	128	0.6	0.5	10.9
Force 192	spiral	120	147	on	on	0.60	192	0.6	0.5	7.2

Name of Series	Thick	Interval	Kernel	Window	IR Lvl	Recon Direction
AX SOFT	5.0	5.0	Br40 / B41f	mediastinum	3	head/feet
COR SOFT	5.0	5.0	Br40 / B41f	mediastinum	3	front/back
SAG SOFT	5.0	5.0	Br40 / B41f	mediastinum	3	left/right
AX 3s	3.0	3.0	Br40 / B41f	mediastinum	3	head/feet

Send the above recons on the pre contrast scan (if without only) or on the post contrast scan (if IV given).

Send only the following recon on the pre contrast scan (if without and with).

AX PRE SOFT	5.0	5.0	Br40 / B41f	mediastinum	3	head/feet
-------------	-----	-----	-------------	-------------	---	-----------

CT Abdomen/Pelvis Routine

GE PARAMETERS & RECONS

	Scan Type	SFOV	kV	mA Range	Noise Index	Smart mA	Slice Thick	Beam Coll	Pitch	Speed	Rot Time	Dose Red	ASIR	Scan Time
LS 16	helical	large	120	50-440	11.57	on	5	20	1.375	27.50	0.6	NA	NA	10.9
Opt 540	helical	large	120	50-440	11.57	on	5	20	1.375	27.50	0.6	NA	NA	10.9
LS VCT 64	helical	large body	120	50-650	11.57	on	5	40	1.375	55.00	0.5	50	50	4.5
Disc VCT 64	helical	large body	120	50-650	11.57	on	5	40	1.375	55.00	0.5	NA	NA	4.5

Name of Series	Thickness	Interval	Recon Algorithm	Window Width/Level	Recon Direction
AX SOFT	5.0	5.0	std full	400/40	head/feet
COR SOFT	5.0	5.0	std full	400/40	front/back
SAG SOFT	5.0	5.0	std full	400/40	left/right
AX 2.5s	2.5	2.5	std full	400/40	head/feet

Must be first recon.

Send the above recons on the pre contrast scan (if without only) or on the post contrast scan (if IV given).

Send only the following recon on the pre contrast scan (if without and with).

AX PRE SOFT	5.0	5.0	std full	400/40	head/feet
-------------	-----	-----	----------	--------	-----------

PHILIPS PARAMETERS & RECONS

	Scan Mode	kV	Avg mAs	Dose Index	3D Dose	Pitch	Detect	Colli	Rot Time	Scan Time
Incisive 128	helical	120	115	20	on	1.00	64	0.625	0.75	9.4

Name of Series	Thick	Interval	Filter	Window	iDose	Recon Direction
AX SOFT	5.0	5.0	B	mediastinum	3	head/feet
COR SOFT	5.0	5.0	B	mediastinum	3	front/back
SAG SOFT	5.0	5.0	B	mediastinum	3	left/right
AX 3s	3.0	3.0	B	mediastinum	3	head/feet

Send the above recons on the pre contrast scan (if without only) or on the post contrast scan (if IV given).

Send only the following recon on the pre contrast scan (if without and with).

AX SOFT PRE	5.0	5.0	B	mediastinum	3	head/feet
-------------	-----	-----	---	-------------	---	-----------